

# TRICO®

## INSTRUCTION

**TRICO, MAPP 18149** - PCS No.: **06520**

**Trico is an oil in water emulsion**

**Contains** 63.96g sheep fat/kg or  
64.6g sheep fat/L

formulated as an oil in water emulsion  
containing 5-chloro-2-methyl-4-isothiazolin-3-one  
and 2-methyl-2H-isothiazol-3-one

P280 Wear protective gloves.

P501 Dispose of contents/container to a licensed  
hazardous waste disposal contractor or  
collection site except for empty clean  
containers which can be disposed of as  
nonhazardous waste.

EUH208 Contains 5-chloro-2-methyl-4-isothiazolin-  
3-one and 2-methyl-2H-isothiazol-3-one.  
May produce an allergic reaction.

EUH401 To avoid risks to human health and the  
environment, comply with the instructions  
for use.

**Read the label before use. Using this product in  
a manner that is inconsistent with the label may  
be an offence. Follow the code of practice for  
using plant protection products.**



## INSTRUCTION

### IMPORTANT INFORMATION. For use only as an animal repellent

#### Crops/situations: Forest

Max. individual dose (l/ha): . . . . . 20      Max. Total Dose (l/ha): . . . . . 20

Annual Max number treatments: . 1      Latest time of application.: . . -

### CROP SPECIFIC INFORMATION AND ENVIRONMENTAL PROTECTION.

#### 1. Control of deer browsing during the winter.

Application rate:            10 - 20 L/ha applied as a spray at a rate of 5 - 10mls/lead shoot.

Number of applications:    1 per annum

Timing of application:      Autumn or early winter before commencement of damage.

The timing of the application is intended to protect the lead shoot in newly established forestry plantations (coniferous and deciduous). Applications in successive years may be necessary depending on the rate of growth of the new plantation.

#### 2. Control of deer browsing during the spring and summer.

Application rate:            10 - 20 L/ha applied as a neat spray at a rate of 5 - 10mls/lead shoot.

Number of applications:    1 per annum

Timing of application:      spring or early summer before commencement of damage. Application should be timed to protect new growth in both coniferous and deciduous plantations.

#### 3. Control of deer fraying.

Application rate:            10 - 20 L/ha applied as a neat spray at a rate of 5 - 10mls/tree. The application should be restricted to the top and centre of the tree.

Number of applications:    1 per annum

Timing of application:      Velvet removal usually occurs in the late summer. The timing should take place when the first signs of damage are observed. Considerable damage can be caused to plantation crops by deer removing velvet from antlers. In addition certain species of deer, particularly roe deer, also use fraying to set visual and olfactory markers.

#### 4. Control of deer damage to bark.

Application rate:            10 - 20 L/ha applied as a neat spray at a rate of 100mls/tree trunk.

Number of applications:    1 per annum

Timing of application:      late autumn to early spring or as soon as the first signs of damage are observed.

Application should be made to the vulnerable part of the trunk (1.0-1.5m above the ground or snow level) at the first signs of damage.

### COMPATIBILITY.

TRICO should not be tank mixed with any other product. DO NOT apply other products either one week before all one week after the application of TRICO

### CONDITIONS OF SUPPLY.

All goods supplied by Kwizda Agro are of high grade and we believe them to be suitable but as we cannot exercise control over their mixing and use, all conditions and warranties, statutory or otherwise, as to the quality of or fitness for any purpose of our goods are excluded and no responsibility will be accepted by us for any danger or injury whatsoever arising from their storage, handling, application or use.

## **SAFETY PRECAUTIONS:**

The following safety precautions are additional to the „H“ and „P“ phrases listed above.

### **Operator protection.**

WHEN USING DO NOT EAT, DRINK OR SMOKE. WASH CONCENTRATE from skin or eyes immediately. DO NOT BREATHE SPRAY. WASH HANDS AND EXPOSED SKIN before eating and drinking and after work. IF YOU FEEL UNWELL, seek medical advice immediately (show the label where possible).

### **Environmental protection.**

DO NOT contaminate water with the product or its container. Do not clean application equipment near surface water. Avoid contamination via drains from farmyards and roads.

The spray should be directed away from water and spray must not be allowed to fall within 1.0m of the top of the bank of a static or flowing water body.

### **Storage and disposal.**

KEEP AWAY FROM FOOD, DRINK AND ANIMAL FEEDINGSTUFFS. KEEP OUT OF REACH OF CHILDREN. KEEP IN ORIGINAL CONTAINER tightly closed, in a safe place. RINSE CONTAINER THOROUGHLY by using an integrated pressure rinsing device if available or manually rinse three times. Add washings into the spray tank at the time of filling and dispose of safely. Disposal of the container should be via a licensed contractor. PROTECT FROM FROST.

### **DIRECTIONS FOR USE.**

IMPORTANT. This information is approved as part of the product label. All instructions within this section must be read carefully in order to obtain safe and successful use of this product.

## **RESTRICTIONS/WARNINGS.**

### **Warnings.**

TRICO is a vertebrate repellent effective by smell therefore avoid contamination or drift onto food or forage crops.

### **Restrictions.**

TRICO should be applied under dry conditions ideally when solar radiation is low to avoid potentially adverse crop effects. If application is made when temperatures are high do not spray onto young shoots directly. If rain occurs during application reapplication will be required in order to maintain the repellent properties of the product.

### **Vertebrate species repelled.**

TRICO is particularly effective against various species of deer (fallow, roe, red and Sika deer). It is not a general vertebrate repellent and is not expected to be effective against foxes, badgers and wild boar.

### **Methods of application.**

TRICO SHOULD BE SHAKEN WELL BEFORE OPENING THE CONTAINER. TRICO is usually applied to forestry trees in the early stages of growth with a knapsack sprayer fitted with a fine solid cone nozzle apart from use against fraying when a flat fan nozzle should be used. TRICO is applied neat without dilution. After application the knapsack sprayer tank, lance and jet should be thoroughly washed out (a minimum of 3 washes with detergent solution) to ensure all TRICO residues are removed from the sprayer.